



## PANDEMIC (H1N1) 2009 VACCINATION PROGRAM Summary for General Practice

### Priority groups

In the initial stages of the rollout of pandemic (H1N1) 2009 vaccination program, which starts 30 September 2009, the emphasis will be on ensuring that people in specified priority groups are offered the vaccine first because they have been identified as being at highest risk from pandemic (H1N1) 2009 influenza. However, vaccination will not be restricted to those identified as being in a priority group. Vaccination of friends, family, and carers of vulnerable people, or anyone who wishes to protect themselves from pandemic influenza is also encouraged. This will help minimise wastage of the vaccine. The priority groups include:

- Children (aged 10 years and over) and adults with underlying chronic medical conditions including:
  - Chronic respiratory conditions, including asthma and COPD
  - Immunosuppression, including HIV/AIDS and use of immunosuppressive drugs
  - Cancer
  - Diabetes mellitus
  - Cardiac disease, excluding simple hypertension
  - Chronic renal disease
  - Chronic metabolic diseases
  - Haemoglobinopathies
  - Chronic neurological diseases
- Individuals with moderate to severe obesity, BMI>35
- Health care and community care workers (including volunteers and students)
- Indigenous people (including Indigenous children aged 10 years and over) and remote and isolated communities with vulnerable people
- Children aged 10 years and over in special schools
- Parents and guardians of children aged 0 to 6 months
- Pregnant women

**The vaccine is available to general practices and other immunisation service providers using the same distribution arrangements as for vaccines in the National Immunisation Program.**

### Dosage and administration

Panvax® H1N1 vaccine is administered by intramuscular or deep subcutaneous injection, usually in the upper arm, as a single dose.

The vaccine is produced by CSL Limited in two sizes of multi-dose vial containing a nominal 5 mL or 10mL of vaccine respectively. **The dosage for adults and children aged 10 years and over is one dose of 15 micrograms in 0.5mL.**

Panvax® H1N1 vaccine has not been approved for children under 10 years of age at this time. Vaccination providers will be notified once the TGA has approved the vaccine for use in children under 10 years of age.

### Using multi-dose vials

Guidelines have been developed by the Australian Technical Advisory Group on Immunisation (ATAGI) and the Royal Australian College of General Practitioners (RACGP) on how to safely handle, store and administer vaccines from multi-dose vials.

A copy of the guidelines will be provided with each vaccine order, and is also available online at [www.healthemergency.gov.au](http://www.healthemergency.gov.au)

### Vaccine safety for pregnant women

Pregnant women are at risk of severe complications if they catch pandemic (H1N1) 2009 influenza, so vaccination is strongly recommended. Similar to seasonal influenza vaccines, Panvax® H1N1 vaccine is an inactivated vaccine and is safe during all stages of pregnancy.

### Contraindications

Panvax® H1N1 vaccine is contraindicated in anyone who has experienced anaphylaxis following a previous dose of any influenza vaccine, or who has experienced anaphylaxis following any vaccine component, including the antibiotics neomycin or polymyxin.

The vaccine should not be used in anyone who has experienced anaphylaxis to eggs or chicken protein.

The Product Information for Panvax® H1N1 vaccine states that immunisation must be postponed in people who have febrile illness or acute infection.

### Precautions

The immunological response may be diminished if the patient is undergoing corticosteroid or immunosuppressant treatment.

**For more information go to [www.healthemergency.gov.au](http://www.healthemergency.gov.au) or contact the Pandemic Hotline: 180 2007  
Adverse events can be reported to TGA in the usual way.**

Patient information sheets are provided with the vaccine for you to hand out to your patients.